

ARG58401 anti-COX7A2 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes COX7A2
Tested Reactivity	Hu
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	COX7A2
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 33-115 of Human COX7A2 (NP_001856.2).
Conjugation	Un-conjugated
Alternate Names	VIIAL; COX7AL; COXVIIa-L; COXVIIAL; Cytochrome c oxidase subunit VIIa-L; Cytochrome c oxidase subunit 7A2, mitochondrial; COX7AL1; Cytochrome c oxidase subunit VIIa-liver/heart; Cytochrome c oxidase subunit VIIaL

Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HeLa	
Observed Size	11 kDa	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	COX7A2
Gene Full Name	cytochrome c oxidase subunit VIIa polypeptide 2 (liver)
Background	Cytochrome c oxidase, the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of three catalytic subunits encoded by mitochondrial genes, and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, while the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes polypeptide 2 (liver isoform) of subunit VIIa, with this polypeptide being present in both muscle and non-muscle tissues. In addition to polypeptide 2, subunit VIIa includes polypeptide 1 (muscle isoform), which is present only in muscle tissues, and a related protein, which is present in all tissues. Alternative splicing results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 4 and 14. [provided by RefSeq, Oct 2009]
Function	This protein is one of the nuclear-coded polypeptide chains of cytochrome c oxidase, the terminal oxidase in mitochondrial electron transport. [UniProt]
Calculated Mw	9 kDa
Cellular Localization	Mitochondrion inner membrane. [UniProt]

Images



HeLa

ARG58401 anti-COX7A2 antibody WB image

Western blot: 25 µg of HeLa cell lysate stained with ARG58401 anti-COX7A2 antibody at 1:1000 dilution.