

ARG58674 anti-EHHADH antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes EHHADH
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	EHHADH
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 444-723 of Human EHHADH (NP_001957.2).
Conjugation	Un-conjugated
Alternate Names	LBP; ECHD; LBFP; L-PBE; PBE; PBFE; EC 1.1.1.35; EC 4.2.1.17; EC 5.3.3.8; FRTS3; Peroxisomal bifunctional enzyme

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	79 kDa	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	EHHADH
Gene Full Name	enoyl-CoA, hydratase/3-hydroxyacyl CoA dehydrogenase
Background	The protein encoded by this gene is a bifunctional enzyme and is one of the four enzymes of the peroxisomal beta-oxidation pathway. The N-terminal region of the encoded protein contains enoyl-CoA hydratase activity while the C-terminal region contains 3-hydroxyacyl-CoA dehydrogenase activity. Defects in this gene are a cause of peroxisomal disorders such as Zellweger syndrome. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2009]
Calculated Mw	79 kDa
PTM	Acetylated, leading to enhanced enzyme activity. Acetylation is enhanced by up to 80% after treatment either with trichostin A (TSA) or with nicotinamide (NAM) with highest increase on Lys-346. Acetylation and enzyme activity increased by about 1.5% on addition of fatty acids. [UniProt]
Cellular Localization	Peroxisome. [UniProt]

Images



ARG58674 anti-EHHADH antibody ICC/IF image

Immunofluorescence: A549 cells stained with ARG58674 anti-EHHADH antibody.