

ARG58818 anti-GluR3 antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes GluR3
Tested Reactivity	Ms, Rat
Predict Reactivity	Hu
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	GluR3
Species	Human
Immunogen	Recombinant protein corresponding to G29-M360 of Human GluR3.
Conjugation	Un-conjugated
Alternate Names	GluA3; GLUR-K3; GluR-3; Glutamate receptor ionotropic, AMPA 3; GLUR3; GluR-K3; MRX94; GLUR-C; GluR-C; AMPA-selective glutamate receptor 3; GLURC; Glutamate receptor 3

Application Instructions

Application table	Application	Dilution
	WB	0.1 - 0.5 µg/ml

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Sodium azide
Stabilizer	5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	GRIA3
Gene Full Name	glutamate receptor, ionotropic, AMPA 3
Background	Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. These receptors are heteromeric protein complexes composed of multiple subunits, arranged to form ligand-gated ion channels. The classification of glutamate receptors is based on their activation by different pharmacologic agonists. The subunit encoded by this gene belongs to a family of AMPA (alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate)-sensitive glutamate receptors, and is subject to RNA editing (AGA->GGA; R->G). Alternative splicing at this locus results in different isoforms, which may vary in their signal transduction properties. [provided by RefSeq, Jul 2008]
Function	Receptor for glutamate that functions as ligand-gated ion channel in the central nervous system and plays an important role in excitatory synaptic transmission. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. In the presence of CACNG4 or CACNG7 or CACNG8, shows resensitization which is characterized by a delayed accumulation of current flux upon continued application of glutamate. [UniProt]
Calculated Mw	101 kDa
PTM	Palmitoylated. Depalmitoylated upon glutamate stimulation. Cys-621 palmitoylation leads to Golgi retention and decreased cell surface expression. In contrast, Cys-847 palmitoylation does not affect cell surface expression but regulates stimulation-dependent endocytosis (By similarity). [UniProt]
Cellular Localization	Cell membrane; Multi-pass membrane protein. Cell junction, synapse, postsynaptic cell membrane; Multi-pass membrane protein. Interaction with CNIH2 and CNIH3 promotes cell surface expression. [UniProt]

Images



ARG58818 anti-GluR3 antibody WB image

Western blot: 50 µg of Rat brain and Mouse brain lysates stained with ARG58818 anti-GluR3 antibody at 0.5 µg/ml dilution.