

ARG58822
anti-HRG / HPRG antibodyPackage: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes HRG / HPRG
Tested Reactivity	Hu
Tested Application	IHC-P
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	HRG / HPRG
Species	Human
Immunogen	Recombinant protein corresponding to V19-V200 of Human HRG.
Conjugation	Un-conjugated
Alternate Names	THPH11; Histidine-rich glycoprotein; Histidine-proline-rich glycoprotein; HPRG; HRGP

Application Instructions

Application table	Application	Dilution
	IHC-P	0.5 - 1 µg/ml

Application Note IHC-P: Antigen Retrieval: By heat mediation.
* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Sodium azide
Stabilizer	5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	HRG
Gene Full Name	histidine-rich glycoprotein
Background	This histidine-rich glycoprotein contains two cystatin-like domains and is located in plasma and platelets. The physiological function has not been determined but it is known that the protein binds heme, dyes and divalent metal ions. The encoded protein also has a peptide that displays antimicrobial activity against <i>C. albicans</i> , <i>E. coli</i> , <i>S. aureus</i> , <i>P. aeruginosa</i> , and <i>E. faecalis</i> . It can inhibit rosette formation and interacts with heparin, thrombospondin and plasminogen. Two of the protein's effects, the inhibition of fibrinolysis and the reduction of inhibition of coagulation, indicate a potential prothrombotic effect. Mutations in this gene lead to thrombophilia due to abnormal histidine-rich glycoprotein levels. [provided by RefSeq, Nov 2014]
Function	Plasma glycoprotein that binds a number of ligands such as heme, heparin, heparan sulfate, thrombospondin, plasminogen, and divalent metal ions. Binds heparin and heparin/glycosaminoglycans in a zinc-dependent manner. Binds heparan sulfate on the surface of liver, lung, kidney and heart endothelial cells. Binds to N-sulfated polysaccharide chains on the surface of liver endothelial cells. Inhibits rosette formation. Acts as an adapter protein and is implicated in regulating many processes such as immune complex and pathogen clearance, cell chemotaxis, cell adhesion, angiogenesis, coagulation and fibrinolysis. Mediates clearance of necrotic cells through enhancing the phagocytosis of necrotic cells in a heparan sulfate-dependent pathway. This process can be regulated by the presence of certain HRG ligands such as heparin and zinc ions. Binds to IgG subclasses of immunoglobins containing kappa and lambda light chains with different affinities regulating their clearance and inhibiting the formation of insoluble immune complexes. Tethers plasminogen to the cell surface. Binds T-cells and alters the cell morphology. Modulates angiogenesis by blocking the CD6-mediated antiangiogenic effect of thrombospondins, THBS1 and THBS2. Acts as a regulator of the vascular endothelial growth factor (VEGF) signaling pathway; inhibits endothelial cell motility by reducing VEGF-induced complex formation between PXN/paxillin and ILK/integrin-linked protein kinase and by promoting inhibition of VEGF-induced tyrosine phosphorylation of focal adhesion kinases and alpha-actinins in endothelial cells. Also plays a role in the regulation of tumor angiogenesis and tumor immune surveillance. Normalizes tumor vessels and promotes antitumor immunity by polarizing tumor-associated macrophages, leading to decreased tumor growth and metastasis. [UniProt]
Calculated Mw	60 kDa
PTM	Proteolytic cleavage produces several HRG fragments which are mostly disulfide-linked and, therefore, not released. Cleavage by plasmin is inhibited in the presence of heparin, zinc ions or in an acidic environment. Cleavage reduces binding of HRG to heparan sulfate, but enhances the ability of HRG to bind and tether plasminogen to the cell surface. On platelet activation, releases a 33 kDa antiangiogenic peptide which encompasses the HRR. Also cleaved in the C-terminal by plasmin. N-glycosylated. [UniProt]
Cellular Localization	Secreted. [UniProt]

Images



ARG58822 anti-HRG / HPRG antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human intestinal cancer stained with ARG58822 anti-HRG / HPRG antibody.