

## ARG59086 anti-RTKN / Rhotekin antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes RTKN / Rhotekin
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	RTKN / Rhotekin
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 254-513 of Human RTKN / Rhotekin (NP_001015056.1).
Conjugation	Un-conjugated
Alternate Names	Rhotekin

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:100
	WB	1:500 - 1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	A431	
Observed Size	65kDa	

### Properties


Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---


Gene Symbol	RTKN
Gene Full Name	rhotekin
Background	This gene encodes a scaffold protein that interacts with GTP-bound Rho proteins. Binding of this protein inhibits the GTPase activity of Rho proteins. This protein may interfere with the conversion of active, GTP-bound Rho to the inactive GDP-bound form by RhoGAP. Rho proteins regulate many important cellular processes, including cytokinesis, transcription, smooth muscle contraction, cell growth and transformation. Dysregulation of the Rho signal transduction pathway has been implicated in many forms of cancer. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]
Function	Mediates Rho signaling to activate NF-kappa-B and may confer increased resistance to apoptosis to cells in gastric tumorigenesis. May play a novel role in the organization of septin structures. [UniProt]
Calculated Mw	63 kDa

## Images



ARG59086 anti-RTKN / Rhotekin antibody ICC/IF image

Immunofluorescence: MCF7 cells stained with ARG59086 anti-RTKN / Rhotekin antibody.



ARG59086 anti-RTKN / Rhotekin antibody WB image

Western blot: 25 µg of A431 cell lysate stained with ARG59086 anti-RTKN / Rhotekin antibody at 1:1000 dilution.