

Product datasheet

info@arigobio.com

ARG59770 anti-HSD17B2 antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes HSD17B2

Tested Reactivity Hu, Ms, Rat

Tested Application WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name HSD17B2

Species Human

Immunogen Recombinant fusion protein corresponding to aa. 118-387 of Human HSD17B2 (NP_002144.1).

Conjugation Un-conjugated

Alternate Names 17-beta-hydroxysteroid dehydrogenase type 2; Short chain dehydrogenase/reductase family 9C

member 2; EC 1.1.1.239; HSD17; Estradiol 17-beta-dehydrogenase 2; 20-alpha-HSD; 17-beta-HSD 2; Microsomal 17-beta-hydroxysteroid dehydrogenase; E2DH; SDR9C2; 20 alpha-hydroxysteroid

dehydrogenase; EC 1.1.1.62; Testosterone 17-beta-dehydrogenase; EDH17B2

Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Mouse liver	
Observed Size	47 kDa	

Properties

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol HSD17B2

Gene Full Name hydroxysteroid (17-beta) dehydrogenase 2


Function Capable of catalyzing the interconversion of testosterone and androstenedione, as well as estradiol and

estrone. Also has 20-alpha-HSD activity. Uses NADH while EDH17B3 uses NADPH. [UniProt]

Calculated Mw 43 kDa

Cellular Localization Membrane; Single-pass type II membrane protein. [UniProt]

Images



ARG59770 anti-HSD17B2 antibody WB image

Western blot: 25 μg of Mouse liver lysate stained with ARG59770 anti-HSD17B2 antibody at 1:1000 dilution.