

## ARG62582 anti-CDKN1C / p57 Kip2 antibody [KP39]

Package: 100 µl  
Store at: -20°C

### Summary

|                     |   |
|---------------------|---|
| Product Description | Mouse Monoclonal antibody [KP39] recognizes CDKN1C / p57 Kip2   |
| Tested Reactivity   | Hu, Ms, Rat   |
| Tested Application  | IHC-P, IP   |
| Host                | Mouse   |
| Clonality           | Monoclonal  |
| Clone               | KP39  |
| Isotype             | IgG2b   |
| Target Name         | CDKN1C / p57 Kip2   |
| Species             | Human   |
| Immunogen           | Recombinant full length protein (Human).  |
| Conjugation         | Un-conjugated   |
| Alternate Names     | Cyclin-dependent kinase inhibitor p57; BWS; WBS; BWCR; KIP2; Cyclin-dependent kinase inhibitor 1C; p57; p57Kip2 |

### Application Instructions

| Application table | Application | Dilution                  |
|-------------------|-------------|---------------------------|
|                   | IHC-P       | 2 - 10 µg/ml              |
|                   | IP          | 2 µg for per mg of lysate |

**Application Note** \* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

### Properties

|                     |  |
|---------------------|--|
| Form                | Liquid   |
| Purification        | IgG purified   |
| Buffer              | 10mM PBS (pH 7.4), 0.2% BSA and 0.09% Sodium azide   |
| Preservative        | 0.09% Sodium azide   |
| Stabilizer          | 0.2% BSA   |
| Concentration       | 0.2 mg/ml  |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |

Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

|                       |  |
|-----------------------|--|
| Database links        | <a href="#">GeneID: 1028 Human</a><br><a href="#">GeneID: 12577 Mouse</a><br><a href="#">Swiss-port # P49918 Human</a><br><a href="#">Swiss-port # P49919 Mouse</a>  |
| Gene Symbol           | CDKN1C   |
| Gene Full Name        | cyclin-dependent kinase inhibitor 1C (p57, Kip2)   |
| Background            | This gene is imprinted, with preferential expression of the maternal allele. The encoded protein is a tight-binding, strong inhibitor of several G1 cyclin/Cdk complexes and a negative regulator of cell proliferation. Mutations in this gene are implicated in sporadic cancers and Beckwith-Wiedemann syndrome, suggesting that this gene is a tumor suppressor candidate. Three transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Oct 2010] |
| Function              | Potent tight-binding inhibitor of several G1 cyclin/CDK complexes (cyclin E-CDK2, cyclin D2-CDK4, and cyclin A-CDK2) and, to lesser extent, of the mitotic cyclin B-CDC2. Negative regulator of cell proliferation. May play a role in maintenance of the non-proliferative state throughout life. [UniProt]   |
| Research Area         | Cancer antibody; Cell Biology and Cellular Response antibody; Gene Regulation antibody   |
| Calculated Mw         | 32 kDa   |
| Cellular Localization | Nucleus  |