

ARG62608 anti-UCHL1 / PGP9.5 antibody [13C4 (I3C4)]

Package: 100 µl, 50 µl
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [13C4 (I3C4)] recognizes UCHL1 / PGP9.5
Tested Reactivity	Hu, Ms, Rat, Dog, Goat, Gpig, Pig, Rb, Sheep, Zfsh
Tested Application	ICC/IF, IHC-Fr, IHC-P, WB
Host	Mouse
Clonality	Monoclonal
Clone	13C4 (I3C4)
Isotype	IgG
Target Name	UCHL1 / PGP9.5
Species	Human
Immunogen	Human brain
Conjugation	Un-conjugated
Alternate Names	PGP95; UCH-L1; PGP9.5; PARK5; Ubiquitin thioesterase L1; HEL-117; Neuron cytoplasmic protein 9.5; Uch-L1; EC 6.-.-.-; PGP 9.5; Ubiquitin carboxyl-terminal hydrolase isozyme L1; NDGOA; EC 3.4.19.12

Application Instructions

Application table	Application	Dilution
	ICC/IF	Assay-dependent
	IHC-Fr	Assay-dependent
	IHC-P	1:30
	WB	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purified Antibody
Buffer	1X PBS and 0.1% Sodium azide
Preservative	0.1% Sodium azide
Concentration	0.2 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.


Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	UCHL1
Gene Full Name	ubiquitin carboxyl-terminal esterase L1 (ubiquitin thiolesterase)
Background	The protein encoded by this gene belongs to the peptidase C12 family. This enzyme is a thiol protease that hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin. This gene is specifically expressed in the neurons and in cells of the diffuse neuroendocrine system. Mutations in this gene may be associated with Parkinson disease.[provided by RefSeq, Sep 2009]
Function	Ubiquitin-protein hydrolase involved both in the processing of ubiquitin precursors and of ubiquitinated proteins. This enzyme is a thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin. Also binds to free monoubiquitin and may prevent its degradation in lysosomes. The homodimer may have ATP-independent ubiquitin ligase activity. [UniProt]
Research Area	Cell Biology and Cellular Response antibody; Gene Regulation antibody; Neuroscience antibody
Calculated Mw	25 kDa
PTM	O-glycosylated.

Images



ARG62608 anti-UCHL1 / PGP9.5 antibody [13C4 (I3C4)] WB image

Western blot: 30 µg of Mouse brain lysate stained with ARG62608 anti-UCHL1 / PGP9.5 antibody [13C4 (I3C4)] at 1:1000 dilution.



ARG62608 anti-UCHL1 / PGP9.5 antibody [13C4 (I3C4)] WB image

Western blot: 30 µg of Rat brain lysate stained with ARG62608 anti-UCHL1 / PGP9.5 antibody [13C4 (I3C4)] at 1:2000 dilution.