

## ARG62725 anti-CD14 antibody [MEM-18]

Package: 100 µg, 50 µg  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody [MEM-18] recognizes CD14
Tested Reactivity	Hu, NHuPrm
Tested Application	ELISA, FACS, FuncSt, IP, WB
Specificity	The clone MEM-18 reacts with CD14, a 53-55 kDa GPI (glycosylphosphatidylinositol)-linked membrane glycoprotein expressed on monocytes, macrophages and weakly on granulocytes; also expressed by most tissue macrophages. In human, the epitope recognized by MEM-18 is located between amino acids 57-64. HLDA III; WS Code M 253 HLDA IV; WS Code M 314 HLDA V; WS Code M MA087 HLDA VI; WS Code M MA95
Host	Mouse
Clonality	Monoclonal
Clone	MEM-18
Isotype	IgG1
Target Name	CD14
Species	Human
Immunogen	A crude mixture of human urinary proteins precipitated by ammonium sulphate from the urine of a patient suffering from proteinuria.
Conjugation	Un-conjugated
Alternate Names	CD antigen CD14; Myeloid cell-specific leucine-rich glycoprotein; Monocyte differentiation antigen CD14

### Application Instructions

Application table	Application	Dilution
	ELISA	Assay-dependent
	FACS	4 µg/ml
	FuncSt	Assay-dependent
	IP	Assay-dependent
	WB	1:100 - 1:1000
Application Note	<p>ELISA: The antibody MEM-18 has been tested as the detection antibody in a sandwich ELISA for analysis of human CD14 in combination with antibody clone [B-A8] (cat. no. ARG62727)</p> <p>Functional study: The antibody MEM-18 completely blocks binding of fluorescein (FITC) - labeled bacterial LPS to the monocyte surface and it also blocks the binding of CD14 to the extracellular TLR2 domain.</p> <p>WB: Use under non-reducing condition.</p> <p>* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations</p>	

should be determined by the scientist.

## Properties


---

Form	Liquid
Purification	Purified by protein A
Purity	> 95% (by SDS-PAGE)
Buffer	PBS (pH 7.4) and 15 mM Sodium azide
Preservative	15 mM Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Database links	<a href="#">GeneID: 929 Human</a> <a href="#">Swiss-port # P08571 Human</a>
Gene Symbol	CD14
Gene Full Name	CD14 molecule
Background	CD14 is a 55 kDa GPI-anchored glycoprotein, constitutively expressed on the surface of mature monocytes, macrophages, and neutrophils, where serves as a multifunctional lipopolysaccharide receptor; it is also released to the serum both as a secreted and enzymatically cleaved GPI-anchored form. CD14 binds lipopolysaccharide molecule in a reaction catalyzed by lipopolysaccharide-binding protein (LBP), an acute phase serum protein. The soluble sCD14 is able to discriminate slight structural differences between lipopolysaccharides and is important for neutralization of serum allochthonous lipopolysaccharides by reconstituted lipoprotein particles. CD14 affects allergic, inflammatory and infectious processes.
Function	In concert with LBP, binds to monomeric lipopolysaccharide and delivers it to the MD-2/TLR4 complex, thereby mediating the innate immune response to bacterial lipopolysaccharide (LPS). Acts via MyD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. Up-regulates cell surface molecules, including adhesion molecules. [UniProt]
Highlight	Related products: <a href="#">CD14 antibodies</a> ; <a href="#">CD14 ELISA Kits</a> ; <a href="#">CD14 Duos / Panels</a> ; <a href="#">Anti-Mouse IgG secondary antibodies</a> ; Related poster download: <a href="#">Toll-like Receptor.pdf</a>
Research Area	Developmental Biology antibody; Immune System antibody; General Lymphocyte Marker Study antibody; Macrophages and neutrophils antibody
Calculated Mw	40 kDa
PTM	N- and O- glycosylated. O-glycosylated with a core 1 or possibly core 8 glycan.



ARG62725 anti-CD14 antibody [MEM-18] WB image

Western blot: 1. Untransfected HEK239 cell lysate, 2. cell culture supernatant from HEK 293 transfected with human CD14 cDNA, 3. HEK 293 cell lysate transfected with human CD14 cDNA stained with ARG62725 anti-CD14 antibody [MEM-18].

---