

ARG62780 anti-CD235a antibody [HIR2]

Package: 100 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [HIR2] recognizes CD235a
Tested Reactivity	Hu
Tested Application	Agg, CyTOF®-candidate, FACS, IHC-Fr, IHC-P
Specificity	The clone HIR2 recognizes N-terminal portion of glycophorin A (and weakly of glycophorin B). Its antigen is expressed on early erythroblasts, late erythroblasts, erythroblasts, mature erythrocytes and the cells of erythroid cell lines K562 and HEL, but not on all other cells. HLDA VII; WS Code 70299
Host	Mouse
Clonality	Monoclonal
Clone	HIR2
Isotype	IgG2b
Target Name	CD235a
Species	Human
Immunogen	Synthetic peptide (Human, N-terminal)
Conjugation	Un-conjugated
Alternate Names	MN; GPERik; MNS; GPA; GPSAT; PAS-2; MN sialoglycoprotein; CD235a; HGpMiV; CD antigen CD235a; HGpMiXI; Sialoglycoprotein alpha; HGpSta(C); Glycophorin-A

Application Instructions

Application table	Application	Dilution
	Agg	Assay-dependent
	CyTOF®-candidate	Assay-dependent
	FACS	1 - 4 µg/ml
	IHC-Fr	Assay-dependent
	IHC-P	10 µg/ml
Application Note	<p>Flow Cytometry: This HIR2 antibody has been tested by flow cytometric analysis of human peripheral blood leukocytes and cell agglutination assay and can be used at approximately 0.1 µg per million cells.</p> <p>Agglutination: The antibody HIR2 agglutinates untreated RBCs but fails to agglutinate papain-treated cells.</p> <p>* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.</p>	


Properties

Form	Liquid
------	--------

Purification	Purified by protein A
Purity	> 95% (by SDS-PAGE)
Buffer	PBS (pH 7.4) and 15 mM Sodium azide
Preservative	15 mM Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Bioinformation

Database links	GeneID: 2993 Human Swiss-port # P02724 Human
Gene Symbol	GYPA
Gene Full Name	glycophorin A (MNS blood group)
Background	CD235a (Glycophorin A, GPA) is a transmembrane sialoglycoprotein expressed on erythrocytes and their precursors. Similarly to glycophorin B (GPB), these molecules provide the cells with a large mucin-like surface, which minimalizes aggregation between erythrocytes in the circulation. GPA is the carrier of blood group M and N specificities, while GPB accounts for S, s and U specificities. CD235a is a receptor of Hsa, an Streptococcus adhesin.
Function	Glycophorin A is the major intrinsic membrane protein of the erythrocyte. The N-terminal glycosylated segment, which lies outside the erythrocyte membrane, has MN blood group receptors. Appears to be important for the function of SLC4A1 and is required for high activity of SLC4A1. May be involved in translocation of SLC4A1 to the plasma membrane. Is a receptor for influenza virus. Is a receptor for Plasmodium falciparum erythrocyte-binding antigen 175 (EBA-175); binding of EBA-175 is dependent on sialic acid residues of the O-linked glycans. Appears to be a receptor for Hepatitis A virus (HAV). [UniProt]
Highlight	Related products: CD235a antibodies ; Anti-Mouse IgG secondary antibodies ; Related news: CyTOF-candidate Antibodies
Research Area	Cell Biology and Cellular Response antibody
Calculated Mw	16 kDa
PTM	The major O-linked glycan are NeuAc-alpha-(2-3)-Gal-beta-(1-3)-[NeuAc-alpha-(2-6)]-GalNAcOH (about 78 %) and NeuAc-alpha-(2-3)-Gal-beta-(1-3)-GalNAcOH (17 %). Minor O-glycans (5 %) include NeuAc-alpha-(2-3)-Gal-beta-(1-3)-[NeuAc-alpha-(2-6)]-GalNAcOH NeuAc-alpha-(2-8)-NeuAc-alpha-(2-3)-Gal-beta-(1-3)-GalNAcOH. About 1% of all O-linked glycans carry blood group A, B and H determinants. They derive from a type-2 precursor core structure, Gal-beta-(1,3)-GlcNAc-beta-1-R, and the antigens are synthesized by addition of fucose (H antigen-specific) and then N-acetylgalactosamine (A antigen-specific) or galactose (B antigen-specific). Specifically O-linked-glycans are NeuAc-alpha-(2-3)-Gal-beta-(1-3)-GalNAcOH-(6-1)-GlcNAc-beta-(4-1)-[Fuc-alpha-(1-2)]-Gal-beta-(3-1)-GalNAc-alpha (about 1%, B antigen-specific) and NeuAc-alpha-(2-3)-Gal-beta-(1-3)-GalNAcOH-(6-1)-GlcNAc-beta-(4-1)-[Fuc-alpha-(1-2)]-Gal-beta (1 %, O antigen-, A antigen- and B antigen-specific).



ARG62780 anti-CD235a antibody [HIR2] IHC-P image

Immunohistochemistry: Paraffin-embedded Human adrenal tissue stained with ARG62780 anti-CD235a antibody [HIR2] at 10 µg/ml dilution.



ARG62780 anti-CD235a antibody [HIR2] FACS image

Flow Cytometry: Human peripheral blood stained with ARG62780 anti-CD235a antibody [HIR2], followed by incubation with APC labelled Goat anti-Mouse secondary antibody.