

# Product datasheet

info@arigobio.com

# ARG63255 anti-SNX18 / SNAG1 antibody

Package: 100 μg Store at: -20°C

#### **Summary**

Product Description Goat Polyclonal antibody recognizes SNX18 / SNAG1

Tested Reactivity Hu
Tested Application WB
Host Goat

**Clonality** Polyclonal

Isotype IgG

Target Name SNX18 / SNAG1

Species Human

Immunogen C-KVQRVPLMTVLSF

Conjugation Un-conjugated

Alternate Names SH3PX2; Sorting nexin-associated Golgi protein 1; Sorting nexin-18; SNAG1; SH3PXD3B; SH3 and PX

domain-containing protein 3B

## **Application Instructions**

Application table	Application	Dilution
	WB	1 - 3 μg/ml
Application Note	WB: Recommend incubate at RT for 1h.  * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations	

\* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

one and the determined by the science

### **Properties**

Form Liquid

**Purification** Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide.

Buffer Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA

Preservative 0.02% Sodium azide

Stabilizer 0.5% BSA

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Database links GeneID: 112574 Human

Swiss-port # Q96RF0 Human

Background This gene encodes a member of the sorting nexin family. Members of this family contain a phox (PX)

domain, which is a phosphoinositide binding domain, and are involved in intracellular trafficking. This protein does not contain a coiled coil region, like some family members, but contains a SH3 domain. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by

RefSeq, Jul 2008]

Research Area Signaling Transduction antibody

Calculated Mw 69 kDa

# **Images**

250kDa
150kDa
100kDa
Western Blot: A549 lysate (35 μg protein in RIPA buffer) stained with
ARG63255 anti-SNX18 / SNAG1 antibody at 1 μg/ml dilution.

50kDa
37kDa
25kDa
20kDa