

ARG63408
anti-SNX8 antibodyPackage: 100 µg
Store at: -20°C

Summary

| | |
|---------------------|--|
| Product Description | Goat Polyclonal antibody recognizes SNX8 |
| Tested Reactivity | Hu |
| Tested Application | IHC-P |
| Host | Goat |
| Clonality | Polyclonal |
| Isotype | IgG |
| Target Name | SNX8 |
| Species | Human |
| Immunogen | TPPCSPPEDGLCPH |
| Conjugation | Un-conjugated |
| Alternate Names | Sorting nexin-8; Mvp1 |

Application Instructions

| Application table | Application | Dilution |
|-------------------|-------------|-------------|
| | IHC-P | 4 - 8 µg/ml |

Application Note IHC-P: Antigen Retrieval: Steam tissue section in Tris/EDTA buffer (pH 9.0).
* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.


Properties

| | |
|---------------------|--|
| Form | Liquid |
| Purification | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| Buffer | Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA |
| Preservative | 0.02% Sodium azide |
| Stabilizer | 0.5% BSA |
| Concentration | 0.5 mg/ml |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note | For laboratory research only, not for drug, diagnostic or other use. |

Bioinformation

| | |
|----------------|---|
| Database links | GeneID: 29886 Human Swiss-port # Q9Y5X2 Human |
| Gene Symbol | SNX8 |
| Gene Full Name | sorting nexin 8 |
| Function | May be involved in several stages of intracellular trafficking. May play a role in intracellular protein transport from early endosomes to the trans-Golgi network. [UniProt] |
| Research Area | Signaling Transduction antibody |
| Calculated Mw | 53 kDa |

Images



ARG63408 anti-SNX8 antibody IHC-P image

Immunohistochemistry: paraffin embedded Human Kidney. (Steamed antigen retrieval with Tris/EDTA buffer pH 9) stained with ARG63408 anti-SNX8 antibody at 4 µg/ml dilution followed by HRP-staining.