

ARG63785 anti-CLCA1 antibody

Package: 100 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes CLCA1
Tested Reactivity	Hu
Tested Application	WB
Specificity	No cross-reactivity expected with CLCA2 and 3.
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	CLCA1
Species	Human
Immunogen	C-TVTSKTNKDTSK
Conjugation	Un-conjugated
Alternate Names	hCLCA1; CACC1; Calcium-activated chloride channel regulator 1; Calcium-activated chloride channel family member 1; CaCC-1; Calcium-activated chloride channel protein 1; EC 3.4.-.-; GOB5; CACC; CLCRG1; hCaCC-1

Application Instructions

Application table	Application	Dilution
	WB	1 - 3 µg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties


Form	Liquid
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 1179 Human Swiss-port # A8K7I4 Human
Background	This gene encodes a member of the calcium sensitive chloride conductance protein family. To date, all members of this gene family map to the same region on chromosome 1p31-p22 and share a high degree of homology in size, sequence, and predicted structure, but differ significantly in their tissue distributions. The encoded protein is expressed as a precursor protein that is processed into two cell-surface-associated subunits, although the site at which the precursor is cleaved has not been precisely determined. The encoded protein may be involved in mediating calcium-activated chloride conductance in the intestine. [provided by RefSeq, Jul 2008]
Research Area	Signaling Transduction antibody
Calculated Mw	100 kDa
PTM	Glycosylated. The 125-kDa product is autoproteolytically processed by the metalloprotease domain and yields to two cell-surface-associated subunits, a 90-kDa protein and a group of 37-to 41-kDa proteins. The cleavage is necessary for calcium-activated chloride channel (CaCC) activation activity.

Images



ARG63785 anti-CLCA1 antibody WB image

Western Blot: Human Duodenum lysate (35 µg protein in RIPA buffer) stained with ARG63785 anti-CLCA1 antibody at 1 µg/ml dilution.