

## ARG64255 anti-CD11b antibody

Package: 100 µg  
Store at: -20°C

### Summary

|                     |  |
|---------------------|--|
| Product Description | Goat Polyclonal antibody recognizes CD11b  |
| Tested Reactivity   | Hu   |
| Tested Application  | ICC/IF   |
| Specificity         | The immunizing peptide represents part of the extra-cellular domain.   |
| Host                | Goat   |
| Clonality           | Polyclonal   |
| Isotype             | IgG  |
| Target Name         | CD11b  |
| Species             | Human  |
| Immunogen           | C-RNDGEDSYRTQ  |
| Conjugation         | Un-conjugated  |
| Alternate Names     | MAC1A; CR3A; CR-3 alpha chain; Cell surface glycoprotein MAC-1 subunit alpha; Integrin alpha-M; MAC-1; CD11 antigen-like family member B; Leukocyte adhesion receptor MO1; MO1A; SLEB6; Neutrophil adherence receptor; CD antigen CD11b; CD11B |

### Application Instructions

|                   |  |          |
|-------------------|--|----------|
| Application table | Application  | Dilution |
|                   | ICC/IF   | 10 µg/ml |
| Application Note  | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. |          |

### Properties


|                     |  |
|---------------------|--|
| Form                | Liquid   |
| Purification        | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.  |
| Buffer              | Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA  |
| Preservative        | 0.02% Sodium azide   |
| Stabilizer          | 0.5% BSA   |
| Concentration       | 0.5 mg/ml  |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |

Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

|                |   |
|----------------|---|
| Database links | <a href="#">GeneID: 3684 Human</a><br><a href="#">Swiss-port # P11215 Human</a>   |
| Background     | CD11b (integrin alpha M chain): Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form a leukocyte-specific integrin referred to as macrophage receptor 1 ('Mac-1'), or inactivated-C3b (iC3b) receptor 3 ('CR3'). The alpha M beta 2 integrin is important in the adherence of neutrophils and monocytes to stimulated endothelium, and also in the phagocytosis of complement coated particles. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2009]  |
| Function       | CD11b: Integrin ITGAM/ITGB2 is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complement-coated particles and pathogens (PubMed:9558116, PubMed:20008295). It is identical with CR-3, the receptor for the iC3b fragment of the third complement component. It probably recognizes the R-G-D peptide in C3b. Integrin ITGAM/ITGB2 is also a receptor for fibrinogen, factor X and ICAM1. It recognizes P1 and P2 peptides of fibrinogen gamma chain. Regulates neutrophil migration (PubMed:28807980). In association with beta subunit ITGB2/CD18, required for CD177-PRTN3-mediated activation of TNF primed neutrophils (PubMed:21193407). May regulate phagocytosis-induced apoptosis in extravasated neutrophils. May play a role in mast cell development. Required with TYROBP/DAP12 in microglia to control production of microglial superoxide ions which promote the neuronal apoptosis that occurs during brain development. [UniProt] |
| Highlight      | Related products:<br><a href="#">CD11b antibodies</a> ; <a href="#">CD11b Duos / Panels</a> ; <a href="#">Anti-Goat IgG secondary antibodies</a> ;<br>Related news:<br><a href="#">New antibody panels and duos for Tumor immune microenvironment</a><br><a href="#">Anti-SerpinB9 therapy, a new strategy for cancer therapy</a>   |
| Research Area  | MDSC Marker antibody; Myeloid-derived suppressor cell antibody  |
| Calculated Mw  | 127 kDa   |

## Images



ARG64255 anti-CD11b antibody ICC/IF image

Immunofluorescence: Paraformaldehyde fixed U937 cells permeabilized with 0.15% Triton. Cells were stained with ARG64255 anti-CD11b antibody (green) at 10 µg/ml dilution for 1 hour. DAPI (blue) for nuclear staining. Negative control: Unimmunized goat IgG (green) at 10 µg/ml dilution.