

Product datasheet

info@arigobio.com

ARG64337 anti-CADM4 antibody

Package: 100 μg Store at: -20°C

Summary

Product Description Goat Polyclonal antibody recognizes CADM4

Tested Reactivity Hu

Predict Reactivity Ms, Rat

Tested Application IHC-P, WB

Host Goat

Clonality Polyclonal

Isotype IgG

Target Name CADM4
Species Human

ImmunogenC-KDERFQLEEFSConjugationUn-conjugated

Alternate Names IGSF4C; NECL4; Cell adhesion molecule 4; IgSF4C; Necl-4; Nectin-like protein 4; TSLC1-like protein 2;

Immunoglobulin superfamily member 4C; TSLL2; NECL-4; synCAM4

Application Instructions

Application table	Application	Dilution
	IHC-P	2.5 μg/ml
	WB	0.1 - 0.3 μg/ml
Application Note	WB: Recommend incubate at RT for 1h. IHC-P: Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form Liquid

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide.

Buffer Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA

Preservative 0.02% Sodium azide

Stabilizer 0.5% BSA

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated

freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links <u>GeneID: 199731 Human</u>

Swiss-port # Q8NFZ8 Human

Gene Symbol CADM4

Gene Full Name cell adhesion molecule 4

Function Involved in the cell-cell adhesion. Has calcium- and magnesium-independent cell-cell adhesion activity.


May have tumor-suppressor activity. [UniProt]

Research Area Cancer antibody

Calculated Mw 43 kDa


PTM N-glycosylated.

Images




ARG64337 anti-CADM4 antibody WB image

Western Blot: Human Brain (Cerebellum) lysate ($35\mu g$ protein in RIPA buffer). A) ARG64337 anti-CADM4 antibody ($0.1\mu g/ml$) and B) ARG64404 anti-CADM4 antibody ($0.05\mu g/ml$).



ARG64337 anti-CADM4 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human skeletal muscle tissue. Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). The tissue section was stained with ARG64337 anti-CADM4 antibody at 2.5 μ g/ml dilution followed by AP-staining.



ARG64337 anti-CADM4 antibody IHC-P image

Immunohistochemistry: paraffin embedded Human Thymus. (Steamed antigen retrieval with citrate buffer pH 6) stained with ARG64337 anti-CADM4 antibody at 2.5 $\mu g/ml$ dilution followed by AP-staining.