

# Product datasheet

info@arigobio.com

# ARG64615 anti-ABCE1/ RNAse L inhibitor antibody

Package: 100 μg Store at: -20°C

## **Summary**

Product Description Goat Polyclonal antibody recognizes ABCE1/ RNAse L inhibitor

Tested Reactivity Hu

Predict Reactivity Ms, Rat, Cow, Dog

Tested Application WB

Specificity Variants (NP\_002931.2; NP\_001035809.1) encode the same protein.

Host Goat

**Clonality** Polyclonal

Isotype IgG

Target Name ABCE1/ RNAse L inhibitor

Species Human

 Immunogen
 C-KLNSIKDVEQKK

 Conjugation
 Un-conjugated

Alternate Names RNS4I; ABC38; 2'-5'-oligoadenylate-binding protein; OABP; RNASEL1; Ribonuclease 4 inhibitor; RLI; ATP-

binding cassette sub-family E member 1; HuHP68; RNase L inhibitor; RNASELI

### **Application Instructions**

Application table	Application	Dilution
	WB	0.3 - 1 μg/ml
Application Note	WB: Recommend incubate at RT for 1h.  * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations	
	should be determined by the scientist.	

### **Properties**

Form Liquid

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide.

Buffer Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA

Preservative 0.02% Sodium azide

Stabilizer 0.5% BSA

Concentration 0.5 mg/ml

**Storage instruction** For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Database links <u>GeneID: 6059 Human</u>

Swiss-port # P61221 Human

Background The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC)

transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the OABP subfamily. Alternatively referred to as the RNase L inhibitor, this protein functions to block the activity of ribonuclease L. Activation of ribonuclease L leads to inhibition of protein synthesis in the 2-5A/RNase L system, the central pathway for viral interferon action. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul

2008]

Research Area Immune System antibody; Signaling Transduction antibody

Calculated Mw 67 kDa

### **Images**

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

#### ARG64615 anti-ABCE1/ RNAse L inhibitor antibody WB image

Western Blot: A431 lysate (35  $\mu g$  protein in RIPA buffer) stained with ARG64615 anti-ABCE1/ RNAse L inhibitor antibody at 0.3  $\mu g/ml$  dilution.