

Product datasheet

info@arigobio.com

ARG64904 anti-APOL6 antibody

Package: 100 μg Store at: -20°C

Summary

Product Description Goat Polyclonal antibody recognizes APOL6

Tested Reactivity Hu
Tested Application WB
Host Goat

Clonality Polyclonal

Isotype IgG

Target Name APOL6
Species Human

Immunogen C-DDIDKTHKKFTKAN

Conjugation Un-conjugated

Alternate Names Apolipoprotein L6; Apolipoprotein L-VI; APOLVI; ApoL-VI

Application Instructions

Application table	Application	Dilution
	WB	1 - 3 µg/ml

Application Note WB: Recommend incubate at RT for 1h.

* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations

should be determined by the scientist.

Properties

Form Liquid

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide.

Buffer Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA

Preservative 0.02% Sodium azide

Stabilizer 0.5% BSA

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links GeneID: 80830 Human

Swiss-port # Q9BWW8 Human

Background This gene is a member of the apolipoprotein L gene family. The encoded protein is found in the


cytoplasm, where it may affect the movement of lipids or allow the binding of lipids to organelles.

[provided by RefSeq, Jul 2008]

Research Area Cell Biology and Cellular Response antibody; Cell Death antibody; Metabolism antibody

Calculated Mw 38 kDa

Images



ARG64904 anti-APOL6 antibody WB image

Western Blot: Human Skeletal Muscle lysate (35 μg protein in RIPA buffer) stained with ARG64904 anti-APOL6 antibody at 2 $\mu g/ml$ dilution.