

Product datasheet

info@arigobio.com

ARG64948 anti-DRAK2 antibody

Package: 100 μg Store at: -20°C

Summary

Product Description Goat Polyclonal antibody recognizes DRAK2

Tested Reactivity Hu
Predict Reactivity Dog
Tested Application WB

Specificity This antiboyd may cross-react with rat STK17B (GeneID: 170904).

Host Goat

Clonality Polyclonal

Isotype IgG

Target Name DRAK2
Species Human

 Immunogen
 C-HSVRSSEDKTSK

 Conjugation
 Un-conjugated

Alternate Names DRAK2; EC 2.7.11.1; DAP kinase-related apoptosis-inducing protein kinase 2; Serine/threonine-protein

kinase 17B

Application Instructions

Application table	Application	Dilution
	WB	1 - 3 μg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations	
	should be determined by the scientist.	

Properties

Form Liquid

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide.

Buffer Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA

Preservative 0.02% Sodium azide

Stabilizer 0.5% BSA

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

www.arigobio.com arigo.nuts about antibodies 1/2

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links GeneID: 9262 Human

Swiss-port # O94768 Human

Gene Symbol STK17B

Gene Full Name serine/threonine kinase 17b

Function Phosphorylates myosin light chains (By similarity). Acts as a positive regulator of apoptosis. [UniProt]

Research Area Cell Biology and Cellular Response antibody; Cell Death antibody; Signaling Transduction antibody

Calculated Mw 42 kDa

PTM Autophosphorylated.

Images

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

ARG64948 anti-DRAK2 antibody WB image

Western Blot: MOLT4 lysate (35 μg protein in RIPA buffer) stained with ARG64948 anti-DRAK2 antibody at 1 $\mu g/ml$ dilution.