

ARG65417 anti-CD103 / Integrin alpha E antibody [Ber-ACT8]

Package: 100 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [Ber-ACT8] recognizes CD103 / Integrin alpha E
Tested Reactivity	Hu, NHuPrm
Tested Application	CyTOF®-candidate, FACS, IHC-Fr, IP, WB
Specificity	The clone Ber-ACT8 recognizes CD103 (alpha E integrin), a type I transmembrane glycoprotein primarily found on intestinal intraepithelial lymphocytes. HLDA V; WS Code A067
Host	Mouse
Clonality	Monoclonal
Clone	Ber-ACT8
Isotype	IgG1
Target Name	CD103 / Integrin alpha E
Species	Human
Immunogen	HTLV-1 induced human T cell line MAPS16
Conjugation	Un-conjugated
Alternate Names	CD103; HUMINAE; Integrin alpha-E; Mucosal lymphocyte 1 antigen; CD antigen CD103; HML-1 antigen; Integrin alpha-IEL

Application Instructions

Application table	Application	Dilution
	CyTOF®-candidate	Assay-dependent
	FACS	1 - 4 µg/ml
	IHC-Fr	Assay-dependent
	IP	Assay-dependent
	WB	Assay-dependent

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

Form	Liquid
Purification	Purified from cell culture supernatant by protein-A affinity chromatography.
Purity	> 95% (by SDS-PAGE)
Buffer	PBS (pH 7.4) and 15 mM Sodium azide

Preservative	15 mM Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 3682 Human Swiss-port # P38570 Human
Gene Symbol	ITGAE
Gene Full Name	integrin, alpha E (antigen CD103, human mucosal lymphocyte antigen 1; alpha polypeptide)
Background	CD103 / Integrin alphaE is an integrin subunit that is expressed on intraepithelial lymphocytes, epithelial dendritic cells, lamina propria-derived dendritic cells, a subpopulation of lamina propria T cells, a small subset of peripheral lymphocytes, namely T reg cells, and on activated and TGF-beta stimulated lymphocytes. CD103 is in mature form cleaved into two disulfide-linked chains (C-terminal 150 kDa chain and N-terminal 25 kDa chain). It heterodimerizes with integrin beta7 subunit to form alphaE/beta7 integrin (mucosal lymphocyte 1 antigen), which through binding E-cadherin mediates homing of lymphocytes to the intestinal epithelium, and, in addition to the role in adhesion, may serve as an accessory molecule for intraepithelial lymphocyte activation.
Function	Integrin alpha-E/beta-7 is a receptor for E-cadherin. It mediates adhesion of intra-epithelial T-lymphocytes to epithelial cell monolayers. [UniProt]
Highlight	Related products: CD103 antibodies ; Anti-Mouse IgG secondary antibodies ; Related news: CyTOF-candidate Antibodies
Research Area	Developmental Biology antibody; Immune System antibody; Signaling Transduction antibody
Calculated Mw	130 kDa