

ARG65507 anti-CD105 / Endoglin antibody [MJ7/18]

Package: 100 µg
Store at: -20°C

Summary

Product Description	Rat Monoclonal antibody [MJ7/18] recognizes CD105 / Endoglin
Tested Reactivity	Ms
Tested Application	FACS, IHC-Fr, IP, WB
Host	Rat
Clonality	Monoclonal
Clone	MJ7/18
Isotype	IgG2a, kappa
Target Name	CD105 / Endoglin
Species	Mouse
Immunogen	Inflamed mouse skin
Conjugation	Un-conjugated
Alternate Names	CD antigen CD105; HHT1; Endoglin; ORW1; END

Application Instructions

Application table	Application	Dilution
	FACS	< 1 µg/10 ⁶ cells
	IHC-Fr	Assay-dependent
	IP	Assay-dependent
	WB	Assay-dependent

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.


Properties

Form	Liquid
Buffer	BBS (pH 8.2)
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 13805 Mouse Swiss-port # Q63961 Mouse
Gene Symbol	Eng
Gene Full Name	endoglin
Background	CD105 (Endoglin) is a homodimeric transmembrane glycoprotein serving in presence of TGFbetaR-2 as a receptor for TGFbeta-1 and TGFbeta-3. CD105 is highly expressed on endothelial cells and promotes angiogenesis during wound healing, infarcts and in a wide range of tumours and its gene expression is stimulated by hypoxia. CD105 prevents apoptosis in hypoxic endothelial cells and also antagonises the inhibitory effects of TGFbeta-1 on vascular endothelial cell growth and migration. Normal cellular levels of CD105 are required for formation of new blood vessels.
Function	Major glycoprotein of vascular endothelium. Involved in the regulation of angiogenesis. May play a critical role in the binding of endothelial cells to integrins and/or other RGD receptors. Acts as TGF-beta coreceptor and is involved in the TGF-beta/BMP signaling cascade. Required for GDF2/BMP9 signaling through SMAD1 in endothelial cells. [UniProt]
Research Area	Cancer antibody; Cell Biology and Cellular Response antibody; Controls and Markers antibody; Developmental Biology antibody; Immune System antibody
Calculated Mw	71 kDa

Images



ARG65507 anti-CD105 / Endoglin antibody [MJ7/18] FACS image

Flow Cytometry: BALB/c Mouse endothelioma cell line bEnd.3 was stained with ARG65507 anti-CD105 / Endoglin antibody [MJ7/18] followed by [ARG21704](#) Mouse anti-Rat IgG2a antibody [2A8F4] (PE).