

ARG66462 anti-DNA Ligase 3 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes DNA Ligase 3
Tested Reactivity	Hu, Ms
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	DNA Ligase 3
Species	Human
Immunogen	KLH-conjugated synthetic peptide within the center region of Human DNA Ligase 3.
Conjugation	Un-conjugated
Alternate Names	DNA ligase III; EC 6.5.1.1; Polydeoxyribonucleotide synthase [ATP] 3; LIG2; DNA ligase 3

Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:1000

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties


Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.42% Potassium phosphate (pH 7.3), 0.87% NaCl, 0.01% Sodium azide and 30% Glycerol.
Preservative	0.01% Sodium azide
Stabilizer	30% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	LIG3
-------------	------

Gene Full Name	ligase III, DNA, ATP-dependent
Background	This gene is a member of the DNA ligase family. Each member of this family encodes a protein that catalyzes the joining of DNA ends but they each have a distinct role in DNA metabolism. The protein encoded by this gene is involved in excision repair and is located in both the mitochondria and nucleus, with translation initiation from the upstream start codon allowing for transport to the mitochondria and translation initiation from a downstream start codon allowing for transport to the nucleus. Additionally, alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008]
Function	Isoform 3 functions as heterodimer with DNA-repair protein XRCC1 in the nucleus and can correct defective DNA strand-break repair and sister chromatid exchange following treatment with ionizing radiation and alkylating agents. Isoform 1 is targeted to mitochondria, where it functions as DNA ligase in mitochondrial base-excision DNA repair. [UniProt]
Calculated Mw	113 kDa
Cellular Localization	Isoform 1: Mitochondrion. Note=Contains an N-terminal mitochondrial transit peptide. Isoform 2: Mitochondrion. Note=Contains an N-terminal mitochondrial transit peptide. Isoform 3: Nucleus. Note=Lacks the N-terminal mitochondrial transit peptide. Isoform 4: Nucleus. Note=Lacks the N-terminal mitochondrial transit peptide. [UniProt]

Images



ARG66462 anti-DNA Ligase 3 antibody WB image

Western blot: Jurkat, NIH/3T3, MCF7, PC3 and HeLa whole cell lysates stained with ARG66462 anti-DNA Ligase 3 antibody.