

## ARG66515 anti-AMACR / p504S antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody recognizes AMACR / p504S
Tested Reactivity	Hu
Tested Application	IHC-P, WB
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1, kappa
Target Name	AMACR / p504S
Species	Human
Immunogen	Synthetic peptide derived from Human AMACR / p504S.
Conjugation	Un-conjugated
Alternate Names	AMACR; Alpha-Methylacyl-CoA Racemase; P504S; RACE; 2-Methylacyl-CoA Racemase; EC 5.1.99.4; AMACRD; CBAS4; RM

### Application Instructions

Application table	Application	Dilution
	IHC-P	1:100 - 1:500
	WB	1:500 - 1:2000
Application Note	IHC-P: Antigen Retrieval: Tris/EDTA buffer (pH 8.0) was used. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	~ 42 kDa	

### Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS, 0.02% Sodium azide, 50% Glycerol and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol and 0.5% BSA
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.


Note

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	AMACR
Gene Full Name	alpha-methylacyl-CoA racemase
Background	This gene encodes a racemase. The encoded enzyme interconverts pristanoyl-CoA and C27-bile acylCoAs between their (R)- and (S)-stereoisomers. The conversion to the (S)-stereoisomers is necessary for degradation of these substrates by peroxisomal beta-oxidation. Encoded proteins from this locus localize to both mitochondria and peroxisomes. Mutations in this gene may be associated with adult-onset sensorimotor neuropathy, pigmentary retinopathy, and adrenomyeloneuropathy due to defects in bile acid synthesis. Alternatively spliced transcript variants have been described. Read-through transcription also exists between this gene and the upstream neighboring C1QTNF3 (C1q and tumor necrosis factor related protein 3) gene.
Function	Acts only on coenzyme A thioesters, not on free fatty acids, and accepts as substrates a wide range of alpha-methylacyl-CoAs, including pristanoyl-CoA, trihydroxycoprostanoyl-CoA (an intermediate in bile acid synthesis), and arylpropionic acids like the anti-inflammatory drug ibuprofen (2-(4-isobutylphenyl)propionic acid) but neither 3-methyl-branched nor linear-chain acyl-CoAs.
Calculated Mw	42 kDa
PTM	Acetylation
Cellular Localization	Mitochondrion, Peroxisome

## Images




ARG66515 anti-AMACR / p504S antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human prostatic carcinoma (high magnification) stained with ARG66515 anti-AMACR / p504S antibody at 1:200 (4°C, overnight). Antigen Retrieval: Tris/EDTA buffer (pH 8.0) was used.



ARG66515 anti-AMACR / p504S antibody WB image

Western blot: 30 µg of HeLa, HepG2 and LnCap whole cell lysates stained with ARG66515 anti-AMACR / p504S antibody at 1:1000 dilution.



**ARG66515 anti-AMACR / p504S antibody IHC-P image**

Immunohistochemistry: Paraffin-embedded Human prostatic carcinoma stained with ARG66515 anti-AMACR / p504S antibody at 1:200 (4°C, overnight). Antigen Retrieval: Tris/EDTA buffer (pH 8.0) was used.