

## ARG66662 anti-Caspase 8 antibody

Package: 100 µg  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody recognizes Caspase 8
Tested Reactivity	Hu, Ms, Rat
Tested Application	IHC-P, WB
Host	Mouse
Clonality	Monoclonal
Isotype	IgG
Target Name	Caspase 8
Species	Human
Immunogen	Recombinant protein of Human Caspase 8.
Conjugation	Un-conjugated
Alternate Names	Casp-8; FADD-like ICE; EC 3.4.22.61; CAP4; ICE-like apoptotic protease 5; MORT1-associated ced-3 homolog; FLICE; Apoptotic cysteine protease; FADD-homologous ICE/ced-3-like protease; Caspase-8; Apoptotic protease Mch-5; CASP-8; MCH5; ALPS2B; MACH

### Application Instructions

Application table	Application	Dilution
	IHC-P	1:200 - 1:500
	WB	1:1000 - 1:2000
Application Note	IHC-P: Antigen Retrieval: Boil tissue section in Sodium citrate buffer (pH 6.0) for 20 min. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	~ 55 kDa	

### Properties


Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.4), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


---

Gene Symbol	CASP8
Gene Full Name	caspase 8, apoptosis-related cysteine peptidase
Background	<p>This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes composed of a prodomain, a large protease subunit, and a small protease subunit. Activation of caspases requires proteolytic processing at conserved internal aspartic residues to generate a heterodimeric enzyme consisting of the large and small subunits. This protein is involved in the programmed cell death induced by Fas and various apoptotic stimuli. The N-terminal FADD-like death effector domain of this protein suggests that it may interact with Fas-interacting protein FADD. This protein was detected in the insoluble fraction of the affected brain region from Huntington disease patients but not in those from normal controls, which implicated the role in neurodegenerative diseases. Many alternatively spliced transcript variants encoding different isoforms have been described, although not all variants have had their full-length sequences determined. [provided by RefSeq, Jul 2008]</p>
Function	<p>Most upstream protease of the activation cascade of caspases responsible for the TNFRSF6/FAS mediated and TNFRSF1A induced cell death. Binding to the adapter molecule FADD recruits it to either receptor. The resulting aggregate called death-inducing signaling complex (DISC) performs CASP8 proteolytic activation. The active dimeric enzyme is then liberated from the DISC and free to activate downstream apoptotic proteases. Proteolytic fragments of the N-terminal propeptide (termed CAP3, CAP5 and CAP6) are likely retained in the DISC. Cleaves and activates CASP3, CASP4, CASP6, CASP7, CASP9 and CASP10. May participate in the GZMB apoptotic pathways. Cleaves ADPRT. Hydrolyzes the small-molecule substrate, Ac-Asp-Glu-Val-Asp- -AMC. Likely target for the cowpox virus CRMA death inhibitory protein. Isoform 5, isoform 6, isoform 7 and isoform 8 lack the catalytic site and may interfere with the pro-apoptotic activity of the complex. [UniProt]</p>
Highlight	<p>Related news: <a href="#">Ripoptosome &amp; Necrosome antibody panels are launched</a></p>
Calculated Mw	55 kDa
PTM	<p>Generation of the subunits requires association with the death-inducing signaling complex (DISC), whereas additional processing is likely due to the autocatalytic activity of the activated protease. GZMB and CASP10 can be involved in these processing events.</p> <p>Phosphorylation on Ser-387 during mitosis by CDK1 inhibits activation by proteolysis and prevents apoptosis. This phosphorylation occurs in cancer cell lines, as well as in primary breast tissues and lymphocytes. [UniProt]</p>
Cellular Localization	Cytoplasm. [UniProt]




ARG66662 anti-Caspase 8 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Mouse spleen tissue stained with ARG66662 anti-Caspase 8 antibody.




ARG66662 anti-Caspase 8 antibody WB image

Western blot: HeLa, Mouse brain and Rat brain lysates stained with ARG66662 anti-Caspase 8 antibody.




ARG66662 anti-Caspase 8 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human tonsil tissue stained with ARG66662 anti-Caspase 8 antibody at 1:200 dilution, overnight at 4°C. Antigen Retrieval: Boil tissue section in Sodium citrate buffer (pH 6.0) for 20 min. Negative control was used by secondary antibody only.



ARG66662 anti-Caspase 8 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Mouse brain tissue stained with ARG66662 anti-Caspase 8 antibody at 1:200 dilution, overnight at 4°C. Antigen Retrieval: Boil tissue section in Sodium citrate buffer (pH 6.0) for 20 min. Negative control was used by secondary antibody only.



ARG66662 anti-Caspase 8 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Mouse liver tissue stained with ARG66662 anti-Caspase 8 antibody (red) at 1:200 dilution, overnight at 4°C. DAPI (blue) for nuclear staining.

---