

### ARG66917 anti-APC15 antibody

Package: 100 μl Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes APC15
Tested Reactivity	Hu
Predict Reactivity	Ms, Rat
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
lsotype	IgG
Target Name	APC15
Species	Human
Immunogen	Synthetic peptide corresponding to Human APC15.
Conjugation	Un-conjugated
Alternate Names	Anaphase-promoting complex subunit 15; HSPC020; C11orf51; APC15

## **Application Instructions**

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Jurkat	
Observed Size	14 kDa	

# Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.4), 0.02% Sodium azide, 50% Glycerol and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol and 0.5% BSA
Concentration	Batch dependent
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

## Bioinformation

Gene Symbol	ANAPC15
Gene Full Name	anaphase promoting complex subunit 15
Function	Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. In the complex, plays a role in the release of the mitotic checkpoint complex (MCC) from the APC/C: not required for APC/C activity itself, but promotes the turnover of CDC20 and MCC on the APC/C, thereby participating in the responsiveness of the spindle assembly checkpoint. Also required for degradation of CDC20. [UniProt]
Calculated Mw	14 kDa
Cellular Localization	cytosol; nucleoplasm [UniProt]