

## ARG66935 anti-STAT6 antibody [SQab22264]

Package: 100 µl, 50 µl  
Store at: -20°C

### Summary

Product Description	Recombinant Rabbit Monoclonal antibody [SQab22264] recognizes STAT6
Tested Reactivity	Hu
Tested Application	IHC-P
Host	Rabbit
Clonality	Monoclonal
Clone	SQab22264
Isotype	IgG
Target Name	STAT6
Species	Human
Immunogen	A synthetic peptide corresponding to the sequence of human Stat6.
Conjugation	Un-conjugated
Alternate Names	D12S1644; STAT6B; STAT6C; Signal transducer and activator of transcription 6; IL-4-STAT; IL-4 Stat

### Application Instructions

Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>IHC-P</td><td>1:100 - 1:200</td></tr></tbody></table>	Application	Dilution	IHC-P	1:100 - 1:200
Application	Dilution				
IHC-P	1:100 - 1:200				
Application Note	IHC-P: Antigen Retrieval: Heat mediated was performed using Tris/EDTA buffer (pH 9.0). Incubate the samples at RT (18-25°C) for 30 min. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.				
Positive Control	Human solitary fibrous tumor (SFT) tissue				

### Properties

Form	Liquid
Purification	Purification with Protein A.
Buffer	PBS, 0.01% Sodium azide, 40% Glycerol and 0.05%BSA.
Preservative	0.01% Sodium azide
Stabilizer	40% Glycerol and 0.05%BSA
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


## Bioinformation

---

Gene Symbol	STAT6
Gene Full Name	signal transducer and activator of transcription 6, interleukin-4 induced
Background	The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein plays a central role in exerting IL4 mediated biological responses. It is found to induce the expression of BCL2L1/BCL-X(L), which is responsible for the anti-apoptotic activity of IL4. Knockout studies in mice suggested the roles of this gene in differentiation of T helper 2 (Th2) cells, expression of cell surface markers, and class switch of immunoglobulins. Alternative splicing results in multiple transcript variants.[provided by RefSeq, May 2010]
Function	Carries out a dual function: signal transduction and activation of transcription. Involved in IL4/interleukin-4- and IL3/interleukin-3-mediated signaling. [UniProt]
Research Area	Gene Regulation antibody; Signaling Transduction antibody
Calculated Mw	94 kDa
PTM	Tyrosine phosphorylated following stimulation by IL4/interleukin-4 and IL3/interleukin-3 (By similarity). Dephosphorylation on tyrosine residues by PTPN2 negatively regulates the IL4/interleukin-4 mediated signaling.

## Images

---



ARG66935 anti-STAT6 antibody [SQab22264] IHC-P image

Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded sections of Human solitary fibrous tumor (SFT) tissue stained with ARG66935 anti-STAT6 antibody [SQab22264]. Antigen Retrieval: Heat tissue section in Tris-EDTA buffer (pH 9.0).