

ARG67005 anti-TLR3 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes TLR3
Tested Reactivity	Hu, Ms
Predict Reactivity	Rat
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	TLR3
Species	Human
Immunogen	Synthetic peptide within the intracellular domain of Human TLR3.
Conjugation	Un-conjugated
Alternate Names	Toll-like receptor 3; CD antigen CD283; CD283; IIAE2

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:200–1:500
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:1000

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	100 mM Tris Glycine (pH 7.0), 0.025% ProClin 300 and 20% Glycerol.
Preservative	0.025% ProClin 300
Stabilizer	20% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	TLR3
Gene Full Name	toll-like receptor 3
Background	<p>TLR3 Antibody: Toll-like receptors (TLRs) are evolutionarily conserved pattern-recognition molecules resembling the toll proteins that mediate antimicrobial responses in <i>Drosophila</i>. These proteins recognize different microbial products during infection and serve as an important link between the innate and adaptive immune responses. The TLRs act through adaptor molecules such as MyD88 and TIRAP to activate various kinases and transcription factors so the organism can respond to potential infection. TLR3 is known to recognize viral double-stranded (ds) RNA, a molecular pattern associated with viral infection. Recently it has been shown to recognize viruses such as Influenza A and West Nile Virus and can mediate entry of at least West Nile Virus. </p>
Highlight	<p>Related products: TLR3 Antibody antibodies; Anti-Rabbit IgG secondary antibodies; Related poster download: Toll-like Receptor.pdf</p>
Research Area	Immune System antibody; Microbiology and Infectious Disease antibody; Signaling Transduction antibody
Calculated Mw	104 kDa
PTM	<p>Heavily N-glycosylated, except on that part of the surface of the ectodomain that is involved in ligand binding.</p> <p>TLR3 signaling requires a proteolytic cleavage mediated by cathepsins CTSB and CTSH, the cleavage occurs between amino acids 252 and 346. The cleaved form of TLR3 is the predominant form found in endosomes.</p>

Images




ARG67005 anti-TLR3 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human cancer tissue stained with ARG67005 anti-TLR3 antibody at 1:200 dilution.



ARG67005 anti-TLR3 antibody WB image

Western blot: Mouse lung tissue stained with ARG67005 anti-TLR3 antibody at 1:800 dilution, overnight at 4°C.



ARG67005 anti-TLR3 antibody ICC/IF image

Western blot: Jurkat cell stained with ARG67005 anti-TLR3 antibody at 1:400 dilution, overnight at 4°C.



ARG67005 anti-TLR3 antibody WB image

Western blot: K562 cells stained with ARG67005 anti-TLR3 antibody at 1:800 dilution, overnight at 4°C.