

ARG67045 anti-ROR gamma T antibody [SQab30332]

Package: 100 µl
Store at: -20°C

Summary

Product Description	Recombinant rabbit Monoclonal antibody [SQab30332] recognizes ROR gamma T
Tested Reactivity	Hu
Tested Application	IHC-P
Host	Rabbit
Clonality	Monoclonal
Clone	SQab30332
Isotype	IgG
Target Name	ROR gamma T
Species	Human
Immunogen	Recombinant protein of Human ROR gamma T
Conjugation	Un-conjugated
Alternate Names	RORC, RAR Related Orphan Receptor C, NR1F3, RZRG, RORG, TOR, Nuclear Receptor Subfamily 1 Group F Member 3, RAR-Related Orphan Receptor C, Nuclear Receptor ROR-Gamma, Nuclear Receptor RZR-Gamma, Retinoid-Related Orphan Receptor Gamma, Retinoid-Related Orphan Receptor-Gamma, Retinoic Acid-Binding Receptor Gamma, RZR-GAMMA, IMD42

Application Instructions

Application table	Application	Dilution
	IHC-P	1:50 - 1:100
Application Note	The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Tonsil	

Properties


Form	Liquid
Purification	Purification with Protein A.
Buffer	PBS, 0.01% Sodium azide, 40% Glycerol and 0.05%BSA.
Preservative	0.01% Sodium azide
Stabilizer	40% Glycerol and 0.05%BSA
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	RORC
Gene Full Name	RAR Related Orphan Receptor C
Background	The protein encoded by this gene is a DNA-binding transcription factor and is a member of the NR1 subfamily of nuclear hormone receptors. The specific functions of this protein are not known; however, studies of a similar gene in mice have shown that this gene may be essential for lymphoid organogenesis and may play an important regulatory role in thymopoiesis. In addition, studies in mice suggest that the protein encoded by this gene may inhibit the expression of Fas ligand and IL2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]
Function	Nuclear receptor that binds DNA as a monomer to ROR response elements (RORE) containing a single core motif half-site 5'-AGGTCA-3' preceded by a short A-T-rich sequence. Key regulator of cellular differentiation, immunity, peripheral circadian rhythm as well as lipid, steroid, xenobiotics and glucose metabolism. [UniProt]
Calculated Mw	58 kDa
Cellular Localization	Nucleus

Images



ARG67045 anti-ROR gamma T antibody [SQab30332] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded tonsil stained with ARG67045 anti-ROR gamma T antibody [SQab30332].