

ARG67057 anti-BACH1 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes BACH1
Tested Reactivity	Hu
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	BACH1
Species	Human
Immunogen	Peptide from human BACH1. AA range:131-180
Conjugation	Un-conjugated
Alternate Names	Transcription regulator protein BACH1; BTBD24; BTB and CNC homolog 1; BACH-1; HA2303

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:200-1:1000
	IHC-P	1:100-1:300
	WB	1:500-1:2000

Application Note IHC-P: Antigen Retrieval: Heat mediation was performed in Sodium citrate buffer (pH 6.0).
* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Observed Size ~ 82 kDa


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium azide.
Preservative	0.02% Sodium azide.
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	BACH1
Gene Full Name	BTB and CNC homology 1, basic leucine zipper transcription factor 1
Background	This gene encodes a transcription factor that belongs to the cap'n'collar type of basic region leucine zipper factor family (CNC-bZip). The encoded protein contains broad complex, tramtrack, bric-a-brac/poxvirus and zinc finger (BTB/POZ) domains, which is atypical of CNC-bZip family members. These BTB/POZ domains facilitate protein-protein interactions and formation of homo- and/or hetero-oligomers. When this encoded protein forms a heterodimer with MafK, it functions as a repressor of Maf recognition element (MARE) and transcription is repressed. Multiple alternatively spliced transcript variants have been identified for this gene. [provided by RefSeq, May 2009]
Function	Transcriptional regulator that acts as repressor or activator. Binds, in vitro, to NF-E2 binding sites. Play important roles in coordinating transcription activation and repression by MAFK. [UniProt]
Calculated Mw	82 kDa
Cellular Localization	Nucleus. [UniProt]

Images




ARG67057 anti-BACH1 antibody IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human tonsil tissue. The section was then stained with ARG67057 anti-BACH1 antibody at 1:200, overnight.



ARG67057 anti-BACH1 antibody ICC/IF image

Immunofluorescence: HeLa cells were stained with ARG67057 anti-BACH1 antibody.



ARG67057 anti-BACH1 antibody WB image

Western blot: HeLa cell lysates stained with ARG67057 anti-BACH1 antibody.