

## ARG67080 anti-CD63 antibody [SQab30352]

Package: 100 µl

Store at: -20°C

### Summary

Product Description	Recombinant Rabbit Monoclonal antibody [SQab30352] recognizes CD63
Tested Reactivity	Hu
Tested Application	IHC-P
Host	Rabbit
Clonality	Monoclonal
Clone	SQab30352
Isotype	IgG
Target Name	CD63
Species	Human
Immunogen	Recombinant protein of Human CD63.
Conjugation	Un-conjugated
Alternate Names	CD63; CD63 Molecule; TSPAN30; ME491; MLA1; Ocular Melanoma-Associated Antigen; CD63 Antigen (Melanoma 1 Antigen); Melanoma-Associated Antigen ME491; Tetraspanin-30; Granulophysin; CD63 Antigen; Tspan-30; OMA81H; Limp1; Lysosomal-Associated Membrane Protein 3; Lysosome Integral Membrane Protein 1; Melanoma-Associated Antigen MLA1; LAMP-3

### Application Instructions

Application table	Application	Dilution
	IHC-P	1:100-1:200
Application Note	IHC-P: Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0). * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Purification with Protein A.
Buffer	PBS, 0.01% Sodium azide, 40% Glycerol and 0.05% BSA.
Preservative	0.01% Sodium azide
Stabilizer	40% Glycerol and 0.05% BSA
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


## Bioinformation

---

Gene Symbol	CD63
Gene Full Name	CD63 Molecule
Background	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. The encoded protein is a cell surface glycoprotein that is known to complex with integrins. It may function as a blood platelet activation marker. Deficiency of this protein is associated with Hermansky-Pudlak syndrome. Also this gene has been associated with tumor progression. Alternative splicing results in multiple transcript variants encoding different protein isoforms. [provided by RefSeq, Apr 2012]
Function	Functions as a cell surface receptor for TIMP1 and plays a role in the activation of cellular signaling cascades. Plays a role in the activation of ITGB1 and integrin signaling, leading to the activation of AKT, FAK/PTK2 and MAP kinases. Promotes cell survival, reorganization of the actin cytoskeleton, cell adhesion, spreading and migration, via its role in the activation of AKT and FAK/PTK2. Plays a role in VEGFA signaling via its role in regulating the internalization of KDR/VEGFR2. Plays a role in intracellular vesicular transport processes, and is required for normal trafficking of the PMEL luminal domain that is essential for the development and maturation of melanocytes. Plays a role in the adhesion of leukocytes onto endothelial cells via its role in the regulation of SELP trafficking. May play a role in mast cell degranulation in response to Ms4a2/FcεRI stimulation, but not in mast cell degranulation in response to other stimuli. [Uniprot]
Calculated Mw	26 kDa
PTM	Glycoprotein, Lipoprotein, Palmitate. [Uniprot]
Cellular Localization	Cell membrane, Endosome, Lysosome, Membrane, Secreted. [Uniprot]

## Images

---



ARG67080 anti-CD63 antibody [SQab30352] IHC-P image

Immunohistochemistry: Human prostate stained with ARG67080 anti-CD63 antibody [SQab30352].