

ARG70496 Mouse CA9 / Carbonic Anhydrase 9 recombinant protein (His-tagged)

Package: 100 µg

Store at: -20°C

Summary

Product Description	CHO expressed, His-tagged Mouse CA9 / Carbonic Anhydrase 9 recombinant protein.
Tested Application	SDS-PAGE
Target Name	CA9 / Carbonic Anhydrase 9
Species	Mouse
A.A. Sequence	Met1-Asp390
Expression System	CHO
Alternate Names	CA9; Carbonic Anhydrase 9; CAIX; MN; Carbonic Anhydrase IX; Renal Cell Carcinoma-Associated Antigen G250; RCC-Associated Protein G250; RCC-Associated Antigen G250; Carbonate Dehydratase IX; Carbonic Dehydratase; Membrane Antigen MN; P54/58N; CA-IX; PMW1; EC 4.2.1.1; G250

Properties

Form	Powder
Purification	>95% (by SDS-PAGE)
Purification Note	Endotoxin level is less than 0.1 EU/µg of the protein, as determined by the LAL test.
Buffer	PBS (pH 7.4)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.
Storage instruction	For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	CA9
Gene Full Name	Carbonic Anhydrase 9
Background	Carbonic anhydrases (CAs) are a large family of zinc metalloenzymes that catalyze the reversible hydration of carbon dioxide. They participate in a variety of biological processes, including respiration, calcification, acid-base balance, bone resorption, and the formation of aqueous humor, cerebrospinal fluid, saliva, and gastric acid. They show extensive diversity in tissue distribution and in their subcellular localization. CA IX is a transmembrane protein and is one of only two tumor-associated carbonic anhydrase isoenzymes known. It is expressed in all clear-cell renal cell carcinoma, but is not detected in normal kidney or most other normal tissues. It may be involved in cell proliferation and transformation. This gene was mapped to 17q21.2 by fluorescence in situ hybridization, however, radiation hybrid mapping localized it to 9p13-p12. [provided by RefSeq, Jun 2014]
Function	Catalyzes the interconversion between carbon dioxide and water and the dissociated ions of carbonic acid (i.e. bicarbonate and hydrogen ions).[Uniprot]