

# Product datasheet

info@arigobio.com

# ARG81272 Human TIMP-2 ELISA Kit

Package: 96 wells Store at: 4°C

## **Summary**

Product Description ARG81272 Human TIMP-2 ELISA Kit is an Enzyme Immunoassay kit for the quantification of Human

TIMP-2 in serum, plasma or cell culture supernatants.

Tested Reactivity Hu

Tested Application ELISA

Target Name TIMP2

Conjugation HRP

Conjugation Note Substrate: TMB and read at 450 nm.

Sensitivity 15 pg/ml

Sample Type Serum, plasma or cell culture supernatants.

Standard Range 31.25 - 2000 pg/ml

Sample Volume  $100 \ \mu l$ 

Alternate Names TIMP-2; Tissue inhibitor of metalloproteinases 2; CSC-21K; DDC8; Metalloproteinase inhibitor 2

## **Application Instructions**

Assay Time ~ 3.5 hours

### **Properties**

Form 96 well

Storage instruction Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test

reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Gene Symbol TIMP2

Gene Full Name TIMP metallopeptidase inhibitor 2

Background This gene is a member of the TIMP gene family. The proteins encoded by this gene family are natural

inhibitors of the matrix metalloproteinases, a group of peptidases involved in degradation of the extracellular matrix. In addition to an inhibitory role against metalloproteinases, the encoded protein has a unique role among TIMP family members in its ability to directly suppress the proliferation of endothelial cells. As a result, the encoded protein may be critical to the maintenance of tissue homeostasis by suppressing the proliferation of quiescent tissues in response to angiogenic factors, and by inhibiting protease activity in tissues undergoing remodelling of the extracellular matrix. [provided

by RefSeq, Jul 2008]

Function Complexes with metalloproteinases (such as collagenases) and irreversibly inactivates them by binding

to their catalytic zinc cofactor. Known to act on MMP-1, MMP-2, MMP-3, MMP-7, MMP-8, MMP-9,

MMP-10, MMP-13, MMP-14, MMP-15, MMP-16 and MMP-19. [UniProt]

Highlight Related products:

TIMP2 antibodies; TIMP2 ELISA Kits;


Related news:

**Detecting MMPs and their non-ECM substrates** 

New ELISA data calculation tool:
Simplify the ELISA analysis by GainData

PTM The activity of TIMP2 is dependent on the presence of disulfide bonds.

### **Images**



#### ARG81272 Human TIMP-2 ELISA Kit standard curve image

ARG81272 Human TIMP-2 ELISA Kit results of a typical standard run with optical density reading at 450 nm.