

# **Product datasheet**

info@arigobio.com

# ARG81685 Mouse CD30 ELISA Kit

Package: 96 wells Store at: 4°C

## Component

Cat. No.	Component Name	Package	Temp
ARG81685-001	Antibody-coated microplate	8 X 12 strips	4°C. Unused strips should be sealed tightly in the air-tight pouch.
ARG81685-002	Standard	2 X 10 ng/vial	4°C
ARG81685-003	Standard/Sample diluent	30 ml (Ready to use)	4°C
ARG81685-004	Antibody conjugate concentrate (100X)	1 vial (100 μl)	4°C
ARG81685-005	Antibody diluent buffer	12 ml (Ready to use)	4°C
ARG81685-006	HRP-Streptavidin concentrate (100X)	1 vial (100 μl)	4°C
ARG81685-007	HRP-Streptavidin diluent buffer	12 ml (Ready to use)	4°C
ARG81685-008	25X Wash buffer	20 ml	4°C
ARG81685-009	TMB substrate	10 ml (Ready to use)	4°C (Protect from light)
ARG81685-010	STOP solution	10 ml (Ready to use)	4°C
ARG81685-011	Plate sealer	4 strips	Room temperature

### **Summary**

Product Description	ARG81685 Mouse CD30 ELISA Kit is an Enzyme Immunoassay kit for the quantification of Mouse CD30

in serum and cell culture supernatants.

Tested Reactivity Ms

Tested Application ELISA

**Specificity** There is no detectable cross-reactivity with other relevant proteins.

Target Name CD30

Conjugation HRP

Conjugation Note Substrate: TMB and read at 450 nm.

Sensitivity 7.8 pg/ml

Sample Type Serum and cell culture supernatants.

Standard Range 15.6 - 1000 pg/ml

Sample Volume  $100 \ \mu l$ 

Precision Intra-Assay CV: 5.8%; Inter-Assay CV: 7.2%

Alternate Names Tumor necrosis factor receptor superfamily member 8; Ki-1 antigen; CD30; Ki-1; Lymphocyte activation

antigen CD30; CD antigen CD30; D1S166E; CD30L receptor

# **Application Instructions**

Assay Time ~ 5 hours

#### **Properties**

Form 96 well

Storage instruction Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test

reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Gene Symbol TNFRSF8

Gene Full Name tumor necrosis factor receptor superfamily, member 8

Background CD30 is a member of the TNF-receptor superfamily. This receptor is expressed by activated, but not by

resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. This receptor is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this

gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

Function CD30 is a receptor for TNFSF8/CD30L (PubMed:8391931). May play a role in the regulation of cellular


growth and transformation of activated lymphoblasts. Regulates gene expression through activation of

NF-kappa-B (PubMed:8999898). [UniProt]

Highlight Related products:

CD30 antibodies; CD30 ELISA Kits; New ELISA data calculation tool: Simplify the ELISA analysis by GainData

PTM Phosphorylated on serine and tyrosine residues. [UniProt]



## ARG81685 Mouse CD30 ELISA Kit standard curve image

ARG81685 Mouse CD30 ELISA Kit results of a typical standard run with optical density reading at 450 nm.