

## ARG82401 Rat JAM-C ELISA Kit

Package: 96 wells  
Store at: 4°C

### Component

Cat. No.	Component Name	Package	Temp
ARG82401-001	Antibody-coated microplate	8 X 12 strips	4°C. Unused strips should be sealed tightly in the air-tight pouch.
ARG82401-002	Standard	2 X 10 ng/vial	4°C
ARG82401-003	Standard/Sample diluent	30 ml (Ready to use)	4°C
ARG82401-004	Antibody conjugate concentrate (100X)	1 vial (100 µl)	4°C
ARG82401-005	Antibody diluent buffer	12 ml (Ready to use)	4°C
ARG82401-006	HRP-Streptavidin concentrate (100X)	1 vial (100 µl)	4°C
ARG82401-007	HRP-Streptavidin diluent buffer	12 ml (Ready to use)	4°C
ARG82401-008	25X Wash buffer	20 ml	4°C
ARG82401-009	TMB substrate	10 ml (Ready to use)	4°C (Protect from light)
ARG82401-010	STOP solution	10 ml (Ready to use)	4°C
ARG82401-011	Plate sealer	4 strips	Room temperature

### Summary

Product Description	ARG82401 Rat JAM-C ELISA Kit is an Enzyme Immunoassay kit for the quantification of Rat JAM-C in serum, plasma (EDTA, heparin) and cell culture supernatants.
Tested Reactivity	Rat
Tested Application	ELISA
Target Name	JAM-C
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	7.8 pg/ml
Sample Type	Serum, plasma (EDTA, heparin) and cell culture supernatants.
Standard Range	15.6 - 1000 pg/ml
Sample Volume	100 µl
Precision	Intra-Assay CV: 4.9% Inter-Assay CV: 6.6%

Alternate Names JAM-2; JAM-3; Junctional adhesion molecule C; JAM-C; Junctional adhesion molecule 3; JAMC

## Application Instructions

---

Assay Time ~ 5 hours

## Properties

---

Form 96 well

Storage instruction Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Gene Symbol JAM3

Gene Full Name junctional adhesion molecule 3


Background Tight junctions represent one mode of cell-to-cell adhesion in epithelial or endothelial cell sheets, forming continuous seals around cells and serving as a physical barrier to prevent solutes and water from passing freely through the paracellular space. The protein encoded by this immunoglobulin superfamily gene member is localized in the tight junctions between high endothelial cells. Unlike other proteins in this family, the this protein is unable to adhere to leukocyte cell lines and only forms weak homotypic interactions. The encoded protein is a member of the junctional adhesion molecule protein family and acts as a receptor for another member of this family. A mutation in an intron of this gene is associated with hemorrhagic destruction of the brain, subependymal calcification, and congenital cataracts. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Apr 2011]

Function Participates in cell-cell adhesion. It is a counter-receptor for ITGAM, mediating leukocyte-platelet interactions and is involved in the regulation of transepithelial migration of polymorphonuclear neutrophils (PMN). The soluble form is a mediator of angiogenesis. [UniProt]

Highlight Related products:  
[JAM-C antibodies](#); [JAM-C ELISA Kits](#);  
New ELISA data calculation tool:  
[Simplify the ELISA analysis by GainData](#)

PTM Proteolytically cleaved from endothelial cells surface into a soluble form by ADAM10 and ADAM17; the release of soluble JAM3 is increased by proinflammatory factors. [UniProt]

Cellular Localization Cell membrane; Single-pass type I membrane protein. Cell junction, desmosome. Secreted, extracellular space. Note=In epithelial cells, it is expressed at desmosomes but not at tight junctions (PubMed:15194813). Localizes at the cell surface of endothelial cells; treatment of endothelial cells with vascular endothelial growth factor stimulates recruitment of JAM3 to cell-cell contacts (PubMed:15994945). [UniProt]



ARG82401 Rat JAM-C ELISA Kit standard curve image

ARG82401 Rat JAM-C ELISA Kit results of a typical standard run with optical density reading at 450 nm.