

ARG82787 Sheep IL4 ELISA Kit

Package: 96 wells

Store at: 4°C

Summary

Product Description	ARG82787 Sheep IL4 ELISA Kit is an Enzyme Immunoassay kit for the quantification of Sheep IL4 in serum, plasma (EDTA) and cell culture supernatants.
Tested Reactivity	Sheep
Tested Application	ELISA
Specificity	Cross-react with Bovine IL4. May cross-react with Goat IL4.
Target Name	IL4
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	15.6 pg/ml
Sample Type	Serum, plasma (EDTA) and cell culture supernatants.
Standard Range	31.25 - 2000 pg/ml
Sample Volume	100 µl
Alternate Names	BSF-1; IL-4; B-cell stimulatory factor 1; BCGF1; Lymphocyte stimulatory factor 1; BCGF-1; Interleukin-4; Pitracinra; Binetrakin; BSF1

Application Instructions

Assay Time	~ 3.5 hours
------------	-------------

Properties

Form	96 well
Storage instruction	Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	IL4
Gene Full Name	interleukin 4
Background	The protein encoded by this gene is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively

spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]


Function

Participates in at least several B-cell activation processes as well as of other cell types (PubMed:3016727). It is a costimulator of DNA-synthesis. It induces the expression of class II MHC molecules on resting B-cells. It enhances both secretion and cell surface expression of IgE and IgG1. It also regulates the expression of the low affinity Fc receptor for IgE (CD23) on both lymphocytes and monocytes. Positively regulates IL31RA expression in macrophages (By similarity). Stimulates autophagy in dendritic cells by interfering with mTORC1 signaling and through the induction of RUFY4 (By similarity). [UniProt]

Cellular Localization

Secreted. [UniProt]

Images



ARG82787 Sheep IL4 ELISA Kit standard curve image

ARG82787 Sheep IL4 ELISA Kit results of a typical standard run with optical density reading at 450 nm.
