

# Product datasheet

info@arigobio.com

## ARG82820 Monkey RANTES ELISA Kit

Package: 96 wells Store at: 4°C

#### Summary

Product Description ARG82820 Monkey RANTES ELISA Kit is an Enzyme Immunoassay kit for the quantification of Monkey

RANTES in serum, plasma and cell culture supernatants.

Tested Reactivity Mk

Tested Application ELISA

Specificity Cross-Reactivity: Not react with ApoAI, BMP7, CCL2, CRP, FGF acidic, HGF, HSP27, IL1 alpha, IFN

gamma, IGF1, MMP2, PDGF, PLA2G7, serpin E1, TGF beta 1, TLR3, TNF alpha and VEGF.

Target Name RANTES

Conjugation HRP

Conjugation Note Substrate: TMB and read at 450 nm.

Sensitivity 7.5 pg/ml

Sample Type Serum, plasma and cell culture supernatants.

Standard Range 15 - 1000 pg/ml

Sample Volume 100 µl

Precision Intra-Assay CV: 6.0%

Inter-Assay CV: 10.0%

Alternate Names 3-68; C-C motif chemokine 5; SIS-delta; D17S136E; SISd; T cell-specific protein P228; EoCP; Small-

inducible cytokine A5; Eosinophil chemotactic cytokine; TCP228; 4-68; RANTES; SCYA5; eoCP; T-cell-

specific protein RANTES

#### **Application Instructions**

Assay Time ~ 3 hours

#### **Properties**

Form 96 well

Storage instruction Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test

reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Gene Symbol CCL5

Gene Full Name chemokine (C-C motif) ligand 5

Background This gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines

form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes.

www.arigobio.com argo.nuts about antibodies 1/2

The superfamily is divided into four subfamilies based on the arrangement of the N-terminal cysteine residues of the mature peptide. This chemokine, a member of the CC subfamily, functions as a chemoattractant for blood monocytes, memory T helper cells and eosinophils. It causes the release of histamine from basophils and activates eosinophils. This cytokine is one of the major HIV-suppressive factors produced by CD8+ cells. It functions as one of the natural ligands for the chemokine receptor chemokine (C-C motif) receptor 5 (CCR5), and it suppresses in vitro replication of the R5 strains of HIV-1, which use CCR5 as a coreceptor. Alternative splicing results in multiple transcript variants that encode different isoforms. [provided by RefSeq, Jul 2013]

**Function** 

Chemoattractant for blood monocytes, memory T-helper cells and eosinophils. Causes the release of histamine from basophils and activates eosinophils. May activate several chemokine receptors including CCR1, CCR3, CCR4 and CCR5. One of the major HIV-suppressive factors produced by CD8+ T-cells. Recombinant RANTES protein induces a dose-dependent inhibition of different strains of HIV-1, HIV-2, and simian immunodeficiency virus (SIV). The processed form RANTES(3-68) acts as a natural chemotaxis inhibitor and is a more potent inhibitor of HIV-1-infection. The second processed form RANTES(4-68) exhibits reduced chemotactic and HIV-suppressive activity compared with RANTES(1-68) and RANTES(3-68) and is generated by an unidentified enzyme associated with monocytes and neutrophils (PubMed:16791620, PubMed:1380064, PubMed:8525373, PubMed:9516414, PubMed:15923218). May also be an agonist of the G protein-coupled receptor GPR75, stimulating inositol trisphosphate production and calcium mobilization through its activation. Together with GPR75, may play a role in neuron survival through activation of a downstream signaling pathway involving the PI3, Akt and MAP kinases. By activating GPR75 may also play a role in insulin secretion by islet cells (PubMed:23979485). [UniProt]

PTM


N-terminal processed form RANTES(3-68) is produced by proteolytic cleavage, probably by DPP4, after secretion from peripheral blood leukocytes and cultured sarcoma cells.

The identity of the O-linked saccharides at Ser-27 and Ser-28 are not reported in PubMed:1380064. They are assigned by similarity. [UniProt]

**Cellular Localization** 

Secreted. [UniProt]

#### **Images**



#### ARG82820 Monkey RANTES ELISA Kit standard curve image

ARG82820 Monkey RANTES ELISA Kit results of a typical standard run with optical density reading at 450 nm.