

## ARG82951 Human Eotaxin ELISA Kit

Package: 96 wells  
Store at: 4°C

### Summary

Product Description	ARG82951 Human Eotaxin ELISA Kit is an Enzyme Immunoassay kit for the quantification of Human Eotaxin in serum, plasma and cell culture supernatants.
Tested Reactivity	Hu
Tested Application	ELISA
Target Name	Eotaxin
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	2.35 pg/ml
Sample Type	Serum, plasma and cell culture supernatants.
Standard Range	4.69 - 300 pg/ml
Sample Volume	100 µl
Alternate Names	SCYA11; C-C motif chemokine 11; Eosinophil chemotactic protein; Eotaxin; Small-inducible cytokine A11

### Application Instructions

Assay Time ~ 3.5 hours

### Properties

Form	96 well
Storage instruction	Store the kit at 4°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Gene Symbol	CCL11
Gene Full Name	chemokine (C-C motif) ligand 11
Background	This antimicrobial gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of the N-terminal cysteine residues of the mature peptide. This chemokine, a member of the CC subfamily, displays chemotactic activity for eosinophils, but not mononuclear cells or neutrophils. This eosinophil-specific chemokine is thought to be involved in eosinophilic inflammatory diseases such as atopic dermatitis, allergic rhinitis, asthma and parasitic infections. [provided by RefSeq, Sep 2014]
Function	In response to the presence of allergens, this protein directly promotes the accumulation of eosinophils, a prominent feature of allergic inflammatory reactions. Binds to CCR3. [UniProt]

## Highlight

Related products:

[Eotaxin antibodies](#); [Eotaxin ELISA Kits](#);

New ELISA data calculation tool:

[Simplify the ELISA analysis by GainData](#)

## PTM

O-linked glycan consists of a Gal-GalNAc disaccharide which is modified with up to 2 sialic acid residues.