

ARG82965 Mouse CCL2 / MCP1 (High sensitive) ELISA kit

Package: 96 wells

Store at: 4°C

Summary

Product Description	ARG82965 Mouse CCL2 / MCP1 (High sensitive) ELISA kit is an Enzyme Immunoassay kit for the quantification of Mouse CCL2 / MCP1 in serum, plasma and cell culture supernatants.
Tested Reactivity	Ms
Tested Application	ELISA
Target Name	CCL2 / MCP1
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	1.95 pg/ml
Sample Type	Serum, plasma and cell culture supernatants.
Standard Range	3.9 - 250 pg/ml
Sample Volume	100 µl
Alternate Names	MCP1; Monocyte chemotactic and activating factor; MCAF; Monocyte chemotactic protein 1; Monocyte secretory protein JE; HSMCR30; Small-inducible cytokine A2; HC11; SMC-CF; GDCF-2; SCYA2; C-C motif chemokine 2; Monocyte chemoattractant protein 1; MCP-1

Application Instructions

Assay Time ~ 3.5 hours

Properties

Form	96 well
Storage instruction	Store the kit at 4°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	Ccl2
Gene Full Name	chemokine (C-C motif) ligand 2
Background	This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine displays chemotactic activity for monocytes and basophils but not for neutrophils or eosinophils. It has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis and atherosclerosis. It binds to chemokine receptors CCR2 and CCR4. [provided by RefSeq, Jul 2008]

Highlight

Related products:

[MCP1 antibodies](#); [MCP1 ELISA Kits](#); [MCP1 Duos / Panels](#);

Related news:

[HMGB1 in inflammation](#)

[Inflammatory Cytokines](#)

[A new role of fat – factory of circulating miRNAs](#)

[Detecting MMPs and their non-ECM substrates](#)

New ELISA data calculation tool:

[Simplify the ELISA analysis by GainData](#)

Research Area

Cancer kit; Cell Biology and Cellular Response kit; Immune System kit; Metabolism kit

PTM

Processing at the N-terminus can regulate receptor and target cell selectivity. Deletion of the N-terminal residue converts it from an activator of basophil to an eosinophil chemoattractant.