

## ARG82982 Mouse TNF alpha ELISA Kit

Package: 96 wells  
Store at: 4°C

### Summary

Product Description	ARG82982 Mouse TNF alpha ELISA Kit is an Enzyme Immunoassay kit for the quantification of Mouse TNF alpha in serum, plasma, cell culture supernatant samples.
Tested Reactivity	Ms
Tested Application	ELISA
Specificity	Cross-Reactivity: Not reacts with Mouse IL8, IL10, TNF RI, TNF RII proteins; Human TNF alpha, TNF beta, IL10, LIF, RANK, Fas proteins and Rat TNF alpha proteins
Target Name	TNF alpha
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	3.9 pg/ml
Sample Type	Serum, plasma and cell culture supernatants.
Standard Range	7.8 - 500 pg/ml
Sample Volume	100 µl
Alternate Names	Tumor necrosis factor ligand superfamily member 2; DIF; Cachectin; ICD2; ICD1; N-terminal fragment; TNF-a; TNFA; TNFSF2; TNF-alpha; Tumor necrosis factor; NTF

### Application Instructions

Assay Time ~ 4 hours

### Properties

Form	96 well
Storage instruction	Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Gene Symbol	Tnf
Gene Full Name	tumor necrosis factor
Background	Tumor Necrosis Factor Alpha (TNF alpha) is a protein secreted by lipopolysaccharide stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF alpha exists as a multimer of two, three, or five noncovalently linked units, but shows a single 17 kDa band following SDS PAGE under non reducing conditions. TNF alpha is closely related to the 25 kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF alpha causes cytolysis

or cytoskeleton of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells, TNF alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophils and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. TNF alpha is currently being evaluated in treatment of certain cancers and AIDS Related Complex.

#### Function

Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFR. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia, Under certain conditions it can stimulate cell proliferation and induce cell differentiation. Impairs regulatory T-cells (Treg) function in individuals with rheumatoid arthritis via FOXP3 dephosphorylation. Upregulates the expression of protein phosphatase 1 (PP1), which dephosphorylates the key 'Ser-418' residue of FOXP3, thereby inactivating FOXP3 and rendering Treg cells functionally defective (PubMed:23396208). The TNF intracellular domain (ICD) form induces IL12 production in dendritic cells. [UniProt]

#### Highlight

Related products:

[TNF alpha antibodies](#); [TNF alpha ELISA Kits](#); [TNF alpha Duos / Panels](#); [TNF alpha recombinant proteins](#);

Related news:

[HMGB1 in inflammation](#)

[Inflammatory Cytokines](#)

[A new role of fat – factory of circulating miRNAs](#)

[LKB1 deficiency in T cells promotes gut tumors](#)

New ELISA data calculation tool:

[Simplify the ELISA analysis by GainData](#)

#### Research Area

Cancer kit; Cell Biology and Cellular Response kit; Immune System kit; Metabolism kit; Signaling Transduction kit

#### PTM

The soluble form derives from the membrane form by proteolytic processing. The membrane-bound form is further proteolytically processed by SPPL2A or SPPL2B through regulated intramembrane proteolysis producing TNF intracellular domains (ICD1 and ICD2) released in the cytosol and TNF C-domain 1 and C-domain 2 secreted into the extracellular space.

The membrane form, but not the soluble form, is phosphorylated on serine residues.

Dephosphorylation of the membrane form occurs by binding to soluble TNFRSF1A/TNFR1.

O-glycosylated; glycans contain galactose, N-acetylgalactosamine and N-acetylneuraminic acid.