

ARG83036 Human Ferritin ELISA Kit

Package: 96 wells
Store at: 4°C

Summary

Product Description	ARG83036 Human Ferritin ELISA Kit is an Enzyme Immunoassay kit for the quantification of Human Ferritin in serum, plasma (heparin), ascites, urine and cell culture supernatants.
Tested Reactivity	Hu
Tested Application	ELISA
Specificity	Not cross-reacts with: Human Hemoglobin, Bilirubin, CD163, SCARA5. Mouse erritin, Hemoglobin, Bilirubin, CD163, SCARA5.
Target Name	FTL
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	62.5 pg/ml
Sample Type	Serum, plasma (heparin), ascites, urine and cell culture supernatants.
Standard Range	125-8000 pg/ml
Sample Volume	100 µl
Alternate Names	FTL, Ferritin Light Chain, Ferritin L Subunit , NBIA3, Ferritin Light Polypeptide Like 3, Ferritin, Light Polypeptide, Ferritin L-Chai, MGC7199, Neurodegeneration With Brain Iron Accumulation 3, Epididymis Secretory Sperm Binding Protei, L Apoferritin, LFTD

Application Instructions

Assay Time ~ 4 hours

Properties

Form	96 well
Storage instruction	Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.


Bioinformation

Gene Symbol	FTL
Gene Full Name	Ferritin light chain
Background	This gene encodes the light subunit of the ferritin protein. Ferritin is the major intracellular iron storage protein in prokaryotes and eukaryotes. It is composed of 24 subunits of the heavy and light ferritin chains. Variation in ferritin subunit composition may affect the rates of iron uptake and release in different tissues. A major function of ferritin is the storage of iron in a soluble and nontoxic state.

Defects in this light chain ferritin gene are associated with several neurodegenerative diseases and hyperferritinemia-cataract syndrome. This gene has multiple pseudogenes. [provided by RefSeq, Jul 2008]

Function	Stores iron in a soluble, non-toxic, readily available form. Important for iron homeostasis. Iron is taken up in the ferrous form and deposited as ferric hydroxides after oxidation. Also plays a role in delivery of iron to cells. Mediates iron uptake in capsule cells of the developing kidney. [UniProt]
Highlight	Related products: Ferritin antibodies; Ferritin ELISA Kits; New ELISA data calculation tool: Simplify the ELISA analysis by GainData
PTM	Acetylation
Cellular Localization	azurophil granule lumen, cytoplasm, cytosol, extracellular exosome, extracellular region, intracellular ferritin complex, membrane. [UniProt]

Images



ARG83036 Human Ferritin ELISA Kit standard curve image

ARG83036 Human Ferritin ELISA Kit results of standard run with optical density reading at 450 nm