

# Product datasheet

info@arigobio.com

# ARG83487 Guinea Pig Complement C3 ELISA Kit

Package: 96 wells Store at: 4°C

## **Summary**

Product Description ARG83487 Guinea Pig Complement C3 ELISA Kit is an Enzyme Immunoassay kit for the quantification of

Guinea Pig Complement C3 in Plasma and Serum

Tested Reactivity Gpig

Tested Application ELISA

Target Name Complement C3

Conjugation HRP

Conjugation Note Substrate: TMB and read at 450 nm.

Sensitivity 1.5 ng/ml

Sample Type Plasma and Serum

Standard Range 3.125 - 200 ng/ml

Sample Volume  $100 \ \mu l$ 

Alternate Names C3; Complement C3; CPAMD1; ARMD9; C3a; C3b; C3 And PZP-Like Alpha-2-Macroglobulin Domain-

Containing Protein 1; Complement Component C3a; Complement Component C3b; Complement Component 3; C3a Anaphylatoxin; Prepro-C3; Epididymis Secretory Sperm Binding Protein Li 62p;

Acylation-Stimulating Protein Cleavage Product; HEL-S-62p; AHUS5; ASP

### **Application Instructions**

Assay Time 0.5 hour

#### **Properties**

Form 96 well

Storage instruction Store the kit at 4°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test

reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Gene Symbol C3

Gene Full Name complement component 3

Background Complement component C3 plays a central role in the activation of complement system. Its activation

is required for both classical and alternative complement activation pathways. The encoded preproprotein is proteolytically processed to generate alpha and beta subunits that form the mature protein, which is then further processed to generate numerous peptide products. The C3a peptide, also

known as the C3a anaphylatoxin, modulates inflammation and possesses antimicrobial activity. Mutations in this gene are associated with atypical hemolytic uremic syndrome and age-related

macular degeneration in human patients

Function C3 plays a central role in the activation of the complement system. Its processing by C3 convertase is

the central reaction in both classical and alternative complement pathways. After activation C3b can

bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates.

PTM Cleavage on pair of basic residues, Disulfide bond, Glycoprotein, Phosphoprotein, Thioester bond

Cellular Localization Secreted